

Period Lengths for Iterated Functions. (Preliminary Version)

Eric Schmutz
Eric.Jonathan.Schmutz@drexel.edu
Drexel Math Department and University of
Delaware Mathematical Sciences Department.

February 2, 2008

Abstract

Let Ω_n be the n^n -element set consisting of functions that have $[n]$ as both domain and codomain. Since Ω_n is finite, it is clear by the pigeonhole principle that, for any $f \in \Omega_n$, the sequence of compositional iterates

$$f, f^{(2)}, f^{(3)}, f^{(4)} \dots$$

must eventually repeat. Let $\mathbf{T}(f)$ be the period of this eventually periodic sequence of functions, i.e. the least positive integer T such that, for all $m \geq n$,

$$f^{(m+T)} = f^{(m)}.$$

A closely related number $\mathbf{B}(f)$ = the product of the lengths of the cycles of f , has previously been used as an approximation for \mathbf{T} . This paper proves that the average values of these two quantities are quite different. The expected value of \mathbf{T} is

$$\frac{1}{n^n} \sum_{f \in \Omega_n} \mathbf{T}(f) = \exp \left(k_0 \sqrt[3]{\frac{n}{\log^2 n}} (1 + o(1)) \right),$$

where k_0 is a complicated but explicitly defined constant that is approximately 3.36. The expected value of \mathbf{B} is much larger:

$$\frac{1}{n^n} \sum_{f \in \Omega_n} \mathbf{B}(f) = \exp \left(\frac{3}{2} \sqrt[3]{n} (1 + o(1)) \right).$$

1 Introduction

Let Ω_n be the n^n -element set consisting of functions that have $[n]$ as both domain and codomain, and let $f^{(t)}$ denote f composed with itself t times.

Since Ω_n is finite, it is clear (by the pigeonhole principle) that, for any $f \in \Omega_n$, the sequence of compositional iterates

$$f, f^{(2)}, f^{(3)}, f^{(4)} \dots$$

must eventually repeat. Define $\mathbf{T}(f)$ to be the period of this sequence, i.e. the least T such that, for all $m \geq n$,

$$f^{(m+T)} = f^{(m)}.$$

We say $v \in [n]$ is a *cyclic vertex* if there is a t such that $f^{(t)}(v) = v$. The restriction of f to cyclic vertices is a permutation of the cyclic vertices, and the period \mathbf{T} is just the order of this permutation, i.e. the least common multiple of the cycle lengths.

Harris showed that $\mathbf{T}(f) = e^{\frac{1}{8} \log^2 n(1+o(1))}$ for most functions f . To make this precise, let P_n denote the uniform distribution on Ω_n ; $P_n(\{f\}) = n^{-n}$ for all f . Define $a_n = \frac{1}{8} \log^2 n$, let $b_n = \frac{1}{\sqrt{24}} \log^{3/2} n$, and let $\phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt$. Using Erdős and Turán's seminal results[9], Harris proved

Theorem 1 (Harris[16]) *For any fixed x ,*

$$\lim_{n \rightarrow \infty} P_n \left(\frac{\log \mathbf{T} - a_n}{b_n} \leq x \right) = \phi(x).$$

Comment: Harris actually stated his theorem for a closely related random variable $\mathbf{O}(f)$ = the number of distinct functions in the sequence $f, f^{(2)}, f^{(3)}, \dots$. However it is clear from his proof that Theorem 1 holds too. In fact, it is straightforward to verify that, for all $f \in \Omega_n$, $|\mathbf{O}(f) - \mathbf{T}(f)| < n$. Such inequalities have been proved by Dénes [8].

Let $\mathbf{B}(f)$ be the product, with multiplicities, of the lengths of the cycles of f . Obviously $\mathbf{T}(f) \leq \mathbf{B}(f)$ for all f , and for some exceptional functions $\mathbf{B}(f)$ is *much* larger than $\mathbf{T}(f)$. For example, if f is a permutation with $n/2$ cycles of length 2, then $\mathbf{B}(f) = 2^{n/2}$, but $\mathbf{T}(f) = 2$. In fact, the maximum value \mathbf{T} can have is $e^{\sqrt{n \log n}(1+o(1))}$ [20]. However, for most random mappings $f \in \Omega_n$, $\mathbf{B}(f)$ is a reasonably good approximation for $\mathbf{T}(f)$. For example, the following proposition follows easily from results of Arratia and Tavaré [3].

Proposition 1 *Let $\ell_n, n = 1, 2, \dots$ be any sequence of positive numbers. There is a constant $c > 0$ such that, for all sufficiently large n ,*

$$P_n(\log \mathbf{B} - \log \mathbf{T} \geq \ell_n) \leq \frac{c \log n (\log \log n)^2}{\ell_n}.$$

Although $\log \mathbf{B}(f)$ and $\log \mathbf{T}(f)$ are approximately equal for most functions f , the set of exceptional functions is nevertheless large enough so that the expected values of the two random variables \mathbf{B} and \mathbf{T} are quite different. The following theorem will be proved.

Theorem 2 $E_n(\mathbf{B}) = \exp \left(\frac{3}{2} \sqrt[3]{n} + O(n^{1/6}) \right)$.

To state a corresponding theorem for \mathbf{T} , we need to define a constant k_0 . First define $I = \int_0^\infty \log \log \left(\frac{e}{1-e^{-t}} \right) dt$, and define $\beta_0 = \sqrt{8I}$. This constant first appears first in [13] where it is proved that the expected order of a random permutation is $\exp \left(\beta_0 \sqrt{n/\log n} (1 + o(1)) \right)$. Also define

$$k_0 = \frac{3}{2} (3I)^{2/3}. \quad (1)$$

We prove

Theorem 3

$$E_n(\mathbf{T}) = \exp \left(k_0 \sqrt[3]{\frac{n}{\log^2 n}} (1 + o(1)) \right).$$

The paper is organized as follows. Section 2 reviews some relevant graph theory. In particular, we need Renyi's asymptotic approximations for the expected length of the cycle in a random f that is chosen uniformly from those $f \in \Omega_n$ that have one cycle. In section 3, we apply Hansen's deconditioning argument so that estimation of $E_n(\mathbf{B})$ can be reduced to a simpler problem of estimating the coefficients of a particular generating function $E_z(\mathbf{B})$. In section 4, we use a Tauberian theorem of Odlyzko to estimate the coefficients of $E_z(\mathbf{B})$ and complete the proof of Theorem 2. In section 5, estimates for $E_n(\mathbf{T})$ are deduced from earlier work on the expected order of a random permutation.

2 The cycle length for a connected map.

If $f \in \Omega_n$, let D_f be the directed graph with vertex set $[n]$ and a directed edge from v to w if and only if $f(v) = w$. Provided we count loops and cycles of length 2, the weak components of D_f each have exactly one directed cycle. Let $U_n \subset \Omega_n$ be the set of functions for which D_f has exactly one weak component. For $f \in U_n$, let $L_n(f)$ be the length of the unique cycle f has. Let κ_n be the expected value of L_n for a uniform random $f \in U_n$. We will need asymptotic estimates for κ_n . Apparently Renyi did this calculation in [22]; see page 366 of Bollobás [7].

Lemma 4 (*Renyi*)

$$|U_n| = \sum_k \binom{n}{k} (k-1)! k n^{n-1-k} = n^n \sqrt{\frac{\pi}{2n}} (1 + O(\frac{1}{\sqrt{n}})),$$

and

$$\kappa_n = \frac{1}{|U_n|} \sum_k k \binom{n}{k} (k-1)! k n^{n-1-k} = \sqrt{\frac{2n}{\pi}} (1 + O(\frac{1}{\sqrt{n}})).$$

Additional background material on random mappings can be found in [12].

3 Deconditioning

The proof of Theorem 3 relies on a probabilistic technique that Jennie Hansen developed for random mappings in [14]. Let P_n be the uniform probability measure on Ω_n , and let $\Omega = \bigcup_{n=1}^{\infty} \Omega_n$. For any $f \in \Omega$ and any positive integer d , let $\alpha_d(f)$ = the number of d -vertex weak components that the graph D_f has. Given $z \in (0, 1)$, and $A \subseteq \Omega$, define $P_z(A)$ to be the probability that the following procedure selects an element of A :

- Choose independently the numbers $\alpha_1, \alpha_2, \alpha_3, \dots$, where α_d has a Poisson distribution with mean $\lambda_d = \frac{z^d}{d} \left(e^{-d} \sum_{k=0}^{d-1} \frac{d^k}{k!} \right)$.
- Let $\nu = \sum_d d\alpha_d$.
- Pick f uniform randomly from among all $f \in \Omega_\nu$ having α_d components of size $d, d = 1, 2, 3, \dots$

This defines a probability measure P_z on Ω . For any $\Psi : \Omega \rightarrow \mathfrak{R}$, let $E_z(\Psi) = \sum_f \Psi(f)P_z(\{f\})$ denote its P_z expected value, and let $E_n(\Psi) = E_z(\Psi|\nu = n) = \frac{1}{n^n} \sum_{f \in \Omega_n} \mathbf{T}(f)$. Hansen proved the following theorem

Theorem 5 ([14]) *If Ψ is any function on Ω that is determined by the sequence $\alpha_1, \alpha_2, \dots$, then*

$$E_n(\Psi) = \frac{e^n n!}{n^n} [[z^n]] H(z) \frac{E_z(\Psi)}{1-z},$$

where $H(z) = \sum_{m=0}^{\infty} \frac{m^m}{m!} (z/e)^m$ and “ $[[z^n]]$ ” means “the coefficient of z^n in \dots ”.

The point of this construction is that $E_z(\Psi)$ is much easier to compute than $E_n(\Psi)$ because the variables $\alpha_d, d = 1, 2, \dots$ are independent with respect to P_z and not with respect to P_n . It is quite analagous to Shepp and Lloyd’s work on random permutations [24].

For the problem under consideration, we let B_d = the product of the lengths of cycles in d -vertex components (If f has no d -vertex components, then $B_d = 1$). Thus $\mathbf{B} = \prod_{d=1}^{\infty} B_d$ is a product of P_z -independent random variables, and

$$E_z(\mathbf{B}) = \prod_{d=1}^{\infty} E_z(B_d). \tag{2}$$

Evaluating the d ’th term in this product, we have

$$E_z(B_d) = \sum_m P_z(\alpha_d = m) E_z(B_d | \alpha_d = m). \tag{3}$$

Note that

$$E_z(B_d | \alpha_d = m) = \kappa_d^m \tag{4}$$

where κ_d = average length of the cycle for a uniform random connected map on d vertices. Therefore

$$E_z(B_d) = e^{-\lambda_d + \lambda_d \kappa_d}. \quad (5)$$

Combining (5), (2), and Theorem 5, we get

$$E_n(\mathbf{B}) = \frac{n!e^n}{n^n} [[z^n]] H(z) \frac{1}{1-z} \exp\left(\sum_{d=1}^{\infty} \lambda_d (\kappa_d - 1)\right). \quad (6)$$

To simplify notation later, define c_d so that $c_d z^d = \lambda_d (\kappa_d - 1)$, i.e.

$$c_d = \frac{(\kappa_d - 1)}{d} (e^{-d} \sum_{k=0}^{d-1} \frac{d^k}{k!}) = \frac{1}{\sqrt{2\pi d}} + O\left(\frac{1}{d}\right). \quad (7)$$

(Further information of the approximation $e^{-d} \sum_{k=0}^{d-1} \frac{d^k}{k!} = \frac{1}{2} + O\left(\frac{1}{\sqrt{d}}\right)$ can be found in [10].) Let $\mu(m) = [[z^m]] \frac{1}{1-z} \exp\left(\sum_{d=1}^{\infty} c_d z^d\right)$, and recall that, for all m , $[[z^m]] H(z) = \frac{m^m}{m!e^m}$. Then

$$E_n(\mathbf{B}) = \frac{n!e^n}{n^n} \sum_{m=0}^n \mu(m) \frac{(n-m)^{n-m}}{(n-m)!e^{n-m}}. \quad (8)$$

In the next section we derive an asymptotic formula for $\log \mu(n)$, and then use it to estimate the right side of (8).

4 Tauberian theorem

In this section, we derive an asymptotic formula for $\log \mu(n)$. Let $a_n = \mu(n) - \mu(n-1)$, and for $s > 0$, let $F(s) = \sum_n a_n e^{-ns} = \exp\left(\sum_{d=1}^{\infty} c_d e^{-ds}\right)$. Also let $g(s) = \log F(s) = \sum_{d=1}^{\infty} c_d e^{-ds}$. To apply Odlyzko's Tauberian theorem[21], we will need asymptotic estimates for the j 'th derivative $g^{(j)}(s) = \sum_{d=1}^{\infty} (-d)^j c_d e^{-ds}$ for $j = 0, 1, 2, 3$. From (7), we have

$$g^{(j)}(s) = \sum_{d=1}^{\infty} \left(\frac{(-d)^j e^{-ds}}{\sqrt{2\pi d}} + O\left(\frac{d^j e^{-ds}}{d}\right) \right). \quad (9)$$

Hence, as $s \rightarrow 0^+$,

$$\begin{aligned} g^{(j)}(s) &= (-1)^j \int_0^{\infty} \frac{t^j e^{-ts}}{\sqrt{2\pi t}} dt + O\left(\int_0^{\infty} t^{j-1} e^{-ts} dt\right) \\ &= \frac{(-1)^j \Gamma(j + \frac{1}{2})}{\sqrt{2\pi} s^{j + \frac{1}{2}}} + O\left(\frac{1}{s^j}\right). \end{aligned} \quad (10)$$

“Rankin’s method” (see Proposition 1 of [21]) is the very useful observation that, for any $s > 0$,

$$\mu(n) \leq e^{ns+g(s)}. \quad (11)$$

In particular, for $s = \frac{1}{2n^{2/3}}$ and $j = 0$, we can combine (11) and (10) to get

$$\mu(n) \leq \exp\left(\frac{3}{2}n^{1/3} + O(1)\right). \quad (12)$$

Now Odlyzko’s Tauberian theorem can be used to derive a lower bound for $\mu(n)$. The remainder of this paragraph is a straightforward verification of the conditions of Theorem 1 in [21]. Let $s_* = s_*(n)$ be chosen so as to minimize $e^{ns+g(s)}$. By calculus and (10),

$$\frac{1}{2n^{2/3}} = s_*(1 + O(\sqrt{s_*})). \quad (13)$$

It is clear that $s_* = o(1)$. We can “bootstrap” twice to get a better estimate. First, by replacing $O(\sqrt{s_*})$ with $o(1)$ in (13), we get $s_* = O(\frac{1}{n^{2/3}})$. Putting this rough bound back into (13) again yields

$$s_* = \frac{1}{2n^{2/3}}(1 + O(\frac{1}{n^{1/3}})). \quad (14)$$

Now let $A_n = g''(s_*)$. By (10) and (14),

$$A_n = 3n^{5/3}(1 + O(\frac{1}{n^{1/3}})). \quad (15)$$

Then, by (10), (14), and (15), we have

$$\left|g'''(s_*)\right| \sim 15n^{7/3} = o(A_n^{3/2}). \quad (16)$$

Having verified the conditions of Odlyzko’s theorem, we conclude that, for all sufficiently large n ,

$$\mu(n) \geq F(s_*) \exp\left(ns_* - 30s_*A_n^{1/2} - 100\right) = e^{\frac{3}{2}\sqrt[3]{n} + O(n^{1/6})}. \quad (17)$$

By Stirling’s formula,

$$\frac{1}{\sqrt{8\pi m}} < \frac{n^m}{m!e^m} < \frac{1}{\sqrt{2\pi m}}. \quad (18)$$

Finally, putting (18) and (17) and (12) back into (8), we get

Theorem 6 $E_n(\mathbf{B}) = e^{\frac{3}{2}\sqrt[3]{n} + O(n^{1/6})}$.

5 Order

The main goal in this section is the proof of Theorem 3, the estimate for the average period $E_n(\mathbf{T})$. However first, for comparison and perspective, we prove Proposition 1, concerning the typical period, that was stated in the introduction.

Proof: Let $Z(f) =$ denote the number of cyclic vertices f has. By the Law of Total Probability,

$$P_n(\log \mathbf{B} - \log \mathbf{T} > \ell_n) = \sum_m P_n(Z = m)P_n(\log \mathbf{B} - \log \mathbf{T} > \ell_n | Z = m). \quad (19)$$

In the proof of Theorem 8, page 333 of [3], Arratia and Tavaré computed the expected value of $\log \mathbf{B} - \log \mathbf{T}$ given the number of cyclic vertices: $E_n(\log \mathbf{B} - \log \mathbf{T} | Z = m) = O(\log m(\log \log m)^2)$. (Note: their use of the notation P_n is not consistent with the notation in this paper.) Therefore, by Markov's inequality, there is a constant $c > 0$ such that, for all $\ell > 0$,

$$P_n(\log \mathbf{B} - \log \mathbf{T} > \ell | Z = m) \leq \frac{c \log m(\log \log m)^2}{\ell} \leq \frac{c \log n(\log \log n)^2}{\ell}. \quad (20)$$

Putting (20) back to the sum (19), we get the proposition. \square

Let $Sym([n]) \subseteq \Omega_n$ be the $n!$ element set of bijections, and let $M_n = E_n(\mathbf{T} | Sym([n])) = \frac{1}{n!} \sum_{\sigma \in Sym([n])} \mathbf{T}(\sigma)$ be the average of the orders of the permutations of $[n]$. Given the set \mathcal{Z} of cyclic vertices, the restriction of a random f to $Sym(\mathcal{Z})$ is a uniform random permutation of \mathcal{Z} . Hence

$$E_n(\mathbf{T}) = \sum_{m=1}^n P_n(Z = m)M_m. \quad (21)$$

Two helpful theorems area make it possible to estimate this sum. The first is a simple formula for $P_n(Z = m)$ that appears in [17] and is attributed to Rubin and Sitgreaves.

Theorem 7 $P_n(Z = m) = \frac{n!m}{(n-m)!n^{m+1}}$

The second helpful theorem is an estimate for M_m . Using Erdős and Turán's Tauberian theorem[9], Richard Stong proved in [25] that

Theorem 8 $\log M_m = \beta_0 \sqrt{m/\log m} + O(\frac{\sqrt{m} \log \log m}{\log m})$.

With Theorems 7 and 8 available, we can prove prove

Theorem 9 $\log E_n(\mathbf{T}) = k_0 \sqrt[3]{\frac{n}{\log^2 n}}(1 + o(1))$.

Proof: Define $a_0 = \sqrt[3]{3I}$, and let $m_0^* =$ the closest integer to $a_0 \sqrt[3]{\frac{n^2}{\log n}}$. For the lower bound, simply let $m = m_0^*$ in the trivial lower bound $E_n(\mathbf{T}) \geq P_n(Z = m)M_m$. Then, by Theorem 7, Theorem 8, and Stirling's formula, $E_n(\mathbf{T})$ is greater than

$$\begin{aligned} & \exp \left(-\frac{(m_0^*)^2}{2n} + O\left(\frac{(m_0^*)^3}{n^2}\right) + \beta_0 \sqrt{\frac{m_0^*}{\log m_0^*}} + O\left(\frac{\sqrt{m_0^*} \log \log m_0^*}{\log m_0^*}\right) \right) \\ & = \exp \left(\frac{k_0 n^{1/3}}{\log^{2/3} n} + O\left(\frac{n^{1/3} \log \log n}{\log^{7/6} n}\right) \right). \end{aligned}$$

For the upper bound, suppose $\epsilon > 0$ is a fixed but arbitrarily small positive number. Define $\beta_\epsilon = \beta + \epsilon$, and $w_\epsilon(m) = \frac{n!}{(n-m)!n^{m-1}} e^{\beta_\epsilon \sqrt{m/\log m}}$. By

Theorem 8, $M_m \leq e^{\beta_\epsilon \sqrt{m/\log m}}$ for all sufficiently large m . Therefore, for all sufficiently large n ,

$$E_n(\mathbf{T}) \leq n \max_{m \leq n} P_n(Z = m) M_m \leq \max_{m \leq n} w_\epsilon(m). \quad (22)$$

For $6 \leq m \leq n$, let $G_{n,\epsilon}(m) = \log w_\epsilon(m)$. If we write $(n-m)! = \Gamma(n+1-m)$, then $G_{n,\epsilon}(x)$ is twice differentiable for all real numbers x with $6 \leq x \leq n$. Let $\Psi(y) = \frac{\Gamma'(y)}{\Gamma(y)}$ be the logarithmic derivative of the Gamma function so that the first two derivatives of $G_{n,\epsilon}$ are

$$G'_{n,\epsilon}(x) = \Psi(n+1-x) - \log n + \frac{\beta_\epsilon}{2\sqrt{x \log x}} \left(1 - \frac{1}{\log x}\right), \quad (23)$$

and

$$G''_{n,\epsilon}(x) = -\Psi'(n+1-x) + \frac{\beta_\epsilon}{4} \frac{(3 - \log^2 x)}{x^{3/2} \log^{5/2} x}. \quad (24)$$

It is well known [4] that $\Psi'(y) = \sum_{k=0}^{\infty} \frac{1}{(y+k)^2} > 0$. Thus both terms in (24) are negative, and we have $G''_{n,\epsilon}(x) < 0$ for $6 \leq x \leq n$. Let $x_{n,\epsilon}^*$ be the unique solution to $G'_{n,\epsilon}(x) = 0$ at which $G_{n,\epsilon}$ attains its maximum. We need to estimate $x_{n,\epsilon}^*$, and then use that estimate to approximate $G_{n,\epsilon}(x_{n,\epsilon}^*)$.

Define $m_{n,\epsilon}^* = \beta_\epsilon^{2/3} \sqrt[3]{3/8} \frac{n^{2/3}}{(\log n)^{1/3}}$, and let $\delta \in (0, \frac{1}{2})$. In order to prove that

$$(1-\delta)m_{n,\epsilon}^* < x_{n,\epsilon}^* < (1+\delta)m_{n,\epsilon}^*, \quad (25)$$

it suffices to verify that $G'_{n,\epsilon}((1-\delta)m_{n,\epsilon}^*) > 0$ and $G'_{n,\epsilon}((1+\delta)m_{n,\epsilon}^*) < 0$. It is well known [4] that

$$\Psi(y) = \log y + O\left(\frac{1}{y}\right). \quad (26)$$

Putting (26) in (23), we get

$$G'_{n,\epsilon}((1-\delta)m_{n,\epsilon}^*) = \frac{\sqrt[3]{3\beta_\epsilon^2}}{\sqrt{1-\delta} \sqrt[3]{8n \log n}} \left\{ 1 - (1-\delta)^{3/2} + O\left(\frac{\log \log n}{\log n}\right) \right\}. \quad (27)$$

To determine the sign, note that $1 - (1-\delta)^{3/2} > \frac{3}{4}\delta$. Hence $G'_{n,\epsilon}((1-\delta)m_{n,\epsilon}^*) > 0$, and we can even allow $\delta = \delta_n$ to depend on n so long as $\frac{\delta \log n}{\log \log n} \rightarrow \infty$. By similar reasoning $G'_{n,\epsilon}((1+\delta)m_{n,\epsilon}^*) < 0$. Therefore $x_{n,\epsilon}^* = m_{n,\epsilon}^* (1 + O(\frac{(\log \log n)^2}{\log n}))$. But then, by Stirling's formula, $G_{n,\epsilon}(x_{n,\epsilon}^*) = \frac{k_\epsilon n^{1/3}}{\log^{2/3} n} (1 + o(1))$, where

$$k_\epsilon = -\frac{(\beta_\epsilon^{2/3} \sqrt[3]{3/8})^2}{2} + \beta_\epsilon \sqrt{\frac{\beta_\epsilon^{2/3} \sqrt[3]{3/8}}{2/3}}.$$

The theorem now follows from the fact that ϵ was an arbitrarily small positive number, and $\lim_{\epsilon \rightarrow 0^+} k_\epsilon = k_0$. \square

6 Discussion

The calculations in this paper depend heavily on the fact that the probability measure P_n is uniform. A separate paper considers Ewens-type distributions where the probability of a permutation or mapping is weighted according to the number of components it has [5],[15],[23]. I do not know if the results in these papers can be extended to more general independent choice models such as those considered by Jaworski in [18] or the p -mappings of Aldous and Pitman, [1],[2].

References

- [1] David Aldous and Jim Pitman, Invariance principles for non-uniform random mappings and trees. Asymptotic combinatorics with application to mathematical physics (St. Petersburg, 2001), 113–147, NATO Sci. Ser. II Math. Phys. Chem., 77, Kluwer Acad. Publ., Dordrecht, 2002.
- [2] David Aldous, Grégory Miermont, and Jim Pitman, Brownian bridge asymptotics for random p -mappings. *Electron. J. Probab.* **9**, no. **3** (2004), 37–56.
- [3] Richard Arratia and Simon Tavaré, Limit theorems for combinatorial structures via discrete process approximations, *Random Structures and Algorithms* **3** (3) (1992)
- [4] Milton Abramowitz and Irene Stegun, Handbook of mathematical functions with formulas, graphs, and mathematical tables, (1992), ISBN 0-486-61272-4.
- [5] A.D. Barbour and Simon Tavaré, A rate for the Erdős Turán law, *Combinatorics, Probability, and Computing* **3** (1994) 167–176.
- [6] A.D. Barbour, Richard Arratia, and Simon Tavaré, “Logarithmic combinatorial structures: a probabilistic approach”, (2003), ISBN 3-03719-000-0.
- [7] Belá Bollobás, Random Graphs, (1985), ISBN 0-12-111755.
- [8] J. Deñes, Some combinatorial properties of transformations and their connections with the theory of graphs. *J. Combinatorial Theory* **9** (1970) 108–116.
- [9] P. Erdős and P. Turán, On some problems of a statistical group theory III, *Acta Math. Acad. Sci. Hung.* **18** (1967) 309–320.
- [10] Phillipe Flajolet, Peter J. Grabner, Peter Kirschenhofer, Helmut Prodinger, On Ramanujan’s Q Function, *Journal of Computational and Applied Mathematics* **58** (1995) 103–116.
- [11] Phillipe Flajolet, Singularity analysis and asymptotics of Bernoulli sums, *Theoretical Computer Science* **215** (1999) 371–381.
- [12] Philippe Flajolet and Andrew M. Odlyzko, Random mapping statistics, *Lecture Notes in Comput. Sci.* **434** (*Advances in cryptology—EUROCRYPT ’89*) (1990) 329–354.

- [13] William Goh and Eric Schmutz, The Expected Order of a Random Permutation, *Bulletin of the London Mathematical Society* **23** (1) (1991),34–42.
- [14] Jennie C.Hansen, A Functional Central Limit Theorem for Random Mappings, *The Annals of Probability*, **17**, No.1(1989) 317–332.
- [15] Jennie C.Hansen, A functional central limit theorem for the Ewens sampling formula. *J. Appl. Probab.* **27** (1990), no. 1, 28–43.
- [16] Bernard Harris, The asymptotic distribution of the order of elements in symmetric semigroups, *Journal of Combinatorial Theory, A* **15** (1973), 66–74.
- [17] Bernard Harris, Probability distributions related to random mappings. *Ann. Math. Statist.* **31** 1960 1045–1062.
- [18] Jerzy Jaworski, On random mapping (T, P_j) , *Journal of Applied Probability* **21** (1) (1984) 186–191.
- [19] Leo Katz, The probability of indecomposability of a random mapping function, *Annals of Mathematical Statistics* **26** (3) (1955) 512–517.
- [20] Massias, J.-P.; Nicolas, J.-L.; Robin, G., Évaluation asymptotique de l'ordre maximum d'un élément du groupe symétrique. *Acta Arith.***50**, no 3 (1988), 221–242.
- [21] Andrew M. Odlyzko, Explicit Tauberian estimates for functions with positive coefficients,*J. Comput. Appl. Math.* **41** (1992), no. 1-2, 187–197.
- [22] Alfred Rényi, On connected graphs. I. *Magyar Tud. Akad. Mat. Kutat Int. Kzl.* **4** 1959 385–388.
- [23] Manuscript in preparation.
- [24] L.A.Shepp and S.P.Lloyd, Ordered cycle lengths in a random permutation. *Trans. Amer. Math. Soc.* **121** 1966 340–357.
- [25] Richard Stong, The average order of a permutation. *Electron. J. Combin.* **5** (1998), Research Paper 41.